

The Circular Chromatic Number of the Mycielski's graph $M^t(K_n)$ *

Zuqiang Ma Junliang Cai

School of Mathematical Sciences, Beijing Normal University

Beijing 100875, P.R.China

(mazuqiang@yahoo.com.cn ; caijunliang@bnu.edu.cn)

Abstract

As a natural generalization of chromatic number of a graph, the circular chromatic number of graphs (or the star chromatic number) was introduced by A.Vince in 1988. Let $M^t(G)$ denote the t th iterated Mycielski graph of G . It was conjectured by Chang, Huang and Zhu (Discrete mathematics, 205(1999), 23-37) that for all $n \geq t + 2$, $\chi_c(M^t(K_n)) = \chi(M^t(K_n)) = n + t$. In 2004, D.D.F. Liu proved the conjecture when $t \geq 2$, $n \geq 2^{t-1} + 2t - 2$. In this paper, we show that the result can be strengthened to the following: if $t \geq 4$, $n \geq \frac{11}{12}2^{t-1} + 2t + \frac{1}{3}$, then $\chi_c(M^t(K_n)) = \chi(M^t(K_n))$.

Keyword: circular chromatic number, complete graph, Mycielski graph

MSC(2000): 05A30

1 Introduction

The circular chromatic number of a graph is a natural generalization of chromatic number $\chi(G)$ of a graph, introduced and studied by A.Vince in 1988, as the ‘star chromatic number’.

Definition 1^[13] Let k and d be positive integers such that $k \geq 2d$, a (k, d) -coloring of a graph G is a mapping

$$f : V(G) \longrightarrow \mathbb{Z}_k = \{0, 1, \dots, k-1\}$$

such that for any $uv \in E(G)$, $d \leq |f(u) - f(v)| \leq k - d$. The circular chromatic number $\chi_c(G)$ of G is defined as

$$\chi_c(G) = \inf \left\{ \frac{k}{d} : \text{there exists a } (k, d)\text{-coloring of } G \right\}.$$

Obviously, a $(k, 1)$ -coloring of a graph G is just an ordinary k -coloring of G . The following properties of circular chromatic number can be found in [2, 13, 14]:

*School of Mathematical Sciences Laboratory of Mathematics and Complex Systems Beijing Normal University, Beijing 100875, China ; Project 10271017 supported by NNSFC

Property 1^[13] $\chi(G) - 1 < \chi_c(G) \leq \chi(G)$.

Property 2^[13] If H is a subgraph of G , then $\chi_c(H) \leq \chi_c(G)$.

Property 3^[2] If there is a homomorphism from G to H , then $\chi_c(G) \leq \chi_c(H)$.

Property 4^[14] For any graph G , $\chi_c(G)$ is a rational number. In fact, we have

$$\chi_c(G) = \min\left\{\frac{k}{d} : \text{there exists a } (k, d)\text{-coloring of } G, k \leq |V(G)|, d \leq \alpha(G)\right\}.$$

where $\alpha(G)$ is the independence number of G .

The question of determining a graph G satisfying $\chi_c(G) = \chi(G)$ was asked by Vince in [13]. Unfortunately it is hard to determine whether a given graph G has $\chi_c(G) = \chi(G)$ or not [5]. In spite of this difficulty, many classes of graphs satisfying $\chi_c(G) = \chi(G)$ have been found, and the Mycielski graph is one of the important classes [3, 4, 10, 11, 12].

Definition 2 Let G be a simple graph with vertex set $V(G) = \{x_1, x_2, \dots, x_n\}$ and edge set $E(G)$, the *Mycielski graph* of G , denoted by $M(G)$, is the graph with vertex set

$$V(M(G)) = \{x_1, x_2, \dots, x_n\} \cup \{x'_1, x'_2, \dots, x'_n\} \cup \{u\},$$

and edge set

$$E(M(G)) = E(G) \cup \{x_i x_j : x_i x_j \in E(G), 1 \leq i, j \leq n\} \cup \{u x_i : 1 \leq i \leq n\}.$$

For each $x_i \in V(G)$, x'_i is called the *twin* of x_i (x_i is also the *twin* of x'_i), and the new vertex u is called the *root* of $M(G)$. Let $M^0(G) = G$, and define the t th Mycielski graph of G is $M^t(G) = M(M^{t-1}(G))$.

Here, we consider $M^i(G)$ as a subgraph of $M^j(G) = M^{j-i}(M^i(G))$ if $i \leq j$.

We know that for any nonempty graph G , $\chi(M(G)) = \chi(G) + 1$. Unfortunately, no simple characterization of graphs with $\chi_c(M(G)) = \chi(M(G))$ have been found, though some graphs that satisfy this condition were studied in [4, 8]. An important class of graphs is the Mycielski graph of complete graph K_n . The following conjecture was introduced by Chang [3] and confirmed for the cases of $t = 1, 2$.

Conjecture^[3] If $t \geq 1, n \geq t + 2$, then $\chi_c(M^t(K_n)) = \chi(M^t(K_n)) = n + t$.

For $t \geq 2$ and $n \geq 2^{t-1} + 2t - 2$, the conjecture is verified [10]. Recently, Simonyi and Tardos proved the conjecture holds when $t + n$ is even.

In this paper, we show that the result in [10] can be strengthened to the following:

Theorem 1 If $t \geq 4, n \geq \frac{11}{12}2^{t-1} + 2t + \frac{1}{3}$, then $\chi_c(M^t(K_n)) = \chi(M^t(K_n)) = n + t$.

2 Some definitions and lemmas

Let $k \geq 2d$, Fan introduced (k, d) -partition of G [4]: a (k, d) -partition of G is a partition $(X_0, X_1, \dots, X_{k-1})$ of $V(G)$ such that $V(G) = \bigcup_{i=0}^{k-1} X_i$, and for each $j \in \mathbb{Z}_{k-1}$,

$$X_j \cup X_{j+1} \cup \dots \cup X_{j+d-1}$$

is an independent set in G (Here it is allowed that $X_i = \emptyset$). It is easy to see that a (k, d) -partition of G is equivalent to a (k, d) -coloring of G (Here we call X_i *color class* of $i, 0 \leq i \leq k-1$). Thus, we have

$$\chi_c(G) = \min\left\{\frac{k}{d} : \text{there exists a } (k, d) \text{-partition of } G\right\}.$$

Combining with the conclusions in [4] and [6], the following result holds:

Lemma 1^[4,6] Let $\chi_c(G) = \frac{k}{d}, (k, d) = 1, (X_0, X_1, \dots, X_{k-1})$ is a (k, d) -partition of G , then $\forall i \in \mathbb{Z}_k, X_i \neq \emptyset$, and $N(X_i) \cap X_{i+d} \neq \emptyset$.

To work with such complicated graphs as $M^t(G)$, we need to take a system to name the vertices of $M^t(G)$.

Definition 3^[11] Suppose $x \in V(G)$, and t is a positive integer,

1. As a vertex of $M^{t-1}(G)$, the twin of x in $M^t(G)$ is called the t th twin of x , denoted by x^t , x is called *initial vertex* of $M^t(G)$;

2. If i and j are positive integers, the i th twin of x in $M^i(G)$ is x^i , the j th of x^i in $M^j(M^i(G)) = M^{i+j}(G)$ is denoted by $(x^i)^j$, which can be simplified as x^{ij} if there is no ambiguity.

If $i_1, i_2, \dots, i_n \in \mathbb{N}$, $x^{i_1 i_2 \dots i_n}$ is defined recursively by $(x^{i_1 i_2 \dots i_{n-1}})^{i_n}$. For any positive integers $i_1, i_2, \dots, i_n$ such that $i_1 + i_2 + \dots + i_n \leq t$, $x^{i_1 i_2 \dots i_n}$ is called a *derived vertex* of x in $M^t(G)$ (Here, the i th twin of x is the derived vertex of x too). The set of derived vertices of x is denoted by $T(x)$:

$$T(x) = \{x^{i_1 i_2 \dots i_n} : i_1 > 0, \dots, i_n > 0, i_1 + i_2 + \dots + i_n \leq t\}.$$

3. For $0 < i \leq t$, the new root in $M^i(G)$ formed from $M^{i-1}(G)$ to $M^i(G)$ is called the i th root of G , denoted by u_i , The set of roots and their derived vertices in $M^t(G)$ is denoted by $R(M^t(G))$:

$$R(M^t(G)) = \{u_i, u_i^{i_1 i_2 \dots i_n} : 0 < i \leq t, i_1 > 0, \dots, i_n > 0, i_1 + i_2 + \dots + i_n \leq t - i\}.$$

Fig.1 shows $M^2(G)$.

For convenience, any vertex in $R(M^t(G))$ is called the *root* of $M^t(G)$.

Definition 4 We call $u_i^{i_1 i_2 \dots i_n}$ the s th *root* of $M^t(G)$ if $i + i_1 + i_2 + \dots + i_n = s$, and denote $R_s(M^t(G))$ as the set of all s th *roots* of $M^t(G)$.

By the definition of Mycielski graph, there is a bijection h from $R(M^{t-1}(G))$ to $R_t(M^t(G)) - \{u_t\}$ as follows

$$h : R(M^{t-1}(G)) \longrightarrow R_t(M^t(G)) - \{u_t\}.$$

For any $u \in R(M^{t-1}(G))$, $h(u)$ is the *twin* of u in $M^t(G) = M(M^{t-1}(G))$. In other words, $h(u_i^{i_1 i_2 \dots i_{n-1}}) = u_i^{i_1 i_2 \dots i_n}$, $i_n \neq 0$, $i_1 + i_2 + \dots + i_n = t$.




Fig.1

We say $x \in V(G)$ is a universal vertex if for each $y \in V(G) - \{x\}$, $xy \in E(G)$. Concerning the circular chromatic number of $M^t(K_n)$, the following results holds:

Lemma 2^[6] Let G be a graph with n universal vertices, $n \geq 2$. If $\chi_c(M^t(G)) = \frac{k}{d}$, $(k, d) = 1$, then $(n - 3)(d - 1) \leq 2^t - 2$.

Lemma 3^[11] Let G be a simple graph, $V(G) = \{x_1, x_2, \dots, x_n\}$. The vertex set of $M(G)$ is $V(M(G)) = \{x_1, x_2, \dots, x_n\} \cup \{x'_1, x'_2, \dots, x'_n\} \cup \{u\}$. If $V(G)$ has (k, d) -partitions, then there is a (k, d) -partition of $V(G)$ such that

1. $u \in X_0$;
2. For some i , $d \leq i \leq k - d$, $\forall x \in V(G)$, we have $\{x, x'\} \subseteq X_i$ if $x \in X_i$.

Definition 5^[11] Let $(X_0, X_1, \dots, X_{k-1})$ be a (k, d) -partition of G . For any $x \in V(G)$, there exist j , $0 \leq j \leq k - 1$ such that $x \in X_j$. The following set

$$\delta(x) = X_{j-d+1} \cup X_{j-d+2} \cup \dots \cup X_j \cup X_{j+1} \cup \dots \cup X_{j+d-1}$$

is called the d -field of x in partition $(X_0, X_1, \dots, X_{k-1})$.

It is easy to see that for any $x \in V(G)$, $\delta(x) \cap N(x) = \emptyset$, where $N(x)$ is the adjacency set of x . In general, we use $C(x)$ to denote the color class containing x .

Suppose $\{x_i, x_j\} \subseteq V(K_n)$, $i \neq j$, by the definition of $M^t(K_n)$, it is obviously that $T(x_i) \subseteq N(x_j)$. Hence, $X_j \subseteq R(M^t(G))$ if $X_j \subseteq \delta(x_i) \cap \delta(x_{i+1})$, because $\forall v \in T(x_l)$, $1 \leq l \leq n$, $v \notin X_j$.

Definition 6 Let $t \geq 1, F_t^\circ$ is a digraph with vertex set

$$V(F_t^\circ) = R_t(M^t(G)) - \{u_t\},$$

and arc set

$$A(F_t^\circ) = \{(u_i^{i_1 i_2 \dots i_n}, u_i^{i_1 i_2 \dots i_{n-1} j(i_n - j)}) : 1 \leq i \leq t-1, i + \sum_{l=1}^n i_l = t, i_n \geq 2, 1 \leq j \leq i_n - 1\}.$$

For convenience of state and proof, we define a directed graph F_t which is the isomorphic graph of F_t° , by replacing $u_i^{i_1 i_2 \dots i_n}$ as $u_i^{i_1 i_2 \dots i_{n-1}}$, $i + i_1 + i_2 + \dots + i_n = t$ (see Fig.2).




Fig.2

Let $F(i), 1 \leq i \leq t-1$, denote the connected component of F_t containing u_i , hence $F_t = \bigcup_{i=1}^{t-1} F(i)$.

Without confusing with the above paragraphs we call u_i the *root* vertex of directed graph $F(i)$ too, $1 \leq i \leq t-1$.

Lemma 4^[7] A weak digraph is an outtree if and only if it has exactly one vertex with indegree 0, and other vertices with indegree 1.

The following lemma is an immediate consequence of Lemma 4 and definition of $F(i)$.

Lemma 5 $\forall i, 1 \leq i \leq t-1$, $F(i)$ is an outtree with *root* u_i , then there is a directed path from u_i to any $u_i^{i_1 i_2 \dots i_n}$.

Let $F'(i)$ be a disjoint isomorphic graph of $F(i)$ with the vertex set $V(F'(i)) = \{v_i^{i_1 i_2 \dots i_n} : i + \sum_{l=1}^n i_l \leq t-1\}$, where $v_i^{i_1 i_2 \dots i_n}$ corresponds to $u_i^{i_1 i_2 \dots i_n} \in V(F(i))$.

Definition 7 For $1 \leq i \leq t-1$, $F(i) \sqcup F'(i)$ is a digraph with vertex set

$$V(F(i) \sqcup F'(i)) = V(F(i)) \cup V(F'(i)),$$

and arc set

$$A(F(i) \sqcup F'(i)) = A(F(i)) \cup A(F'(i)) \cup \{(u_i, v_i)\}.$$

Lemma 6 For $1 \leq i \leq t-2$, $F(i) \cong F(i+1) \sqcup F'(i+1)$.

proof Let g be a mapping from $V(F(i))$ to $V(F(i+1) \sqcup F'(i+1))$, i.e.,

$$g : V(F(i)) \longrightarrow V(F(i+1) \sqcup F'(i+1))$$

such that

$$g(x) = \begin{cases} u_{i+1}, & x = u_i; \\ u_{i+1}^{(i_1-1)i_2 \cdots i_n}, & x = u_i^{i_1 i_2 \cdots i_n}, i_1 \geq 2; \\ v_{i+1}, & x = u_i^1; \\ v_{i+1}^{i_2 \cdots i_n}, & x = u_i^{1 i_2 \cdots i_n}. \end{cases}$$

Then $(u_i, u_i^1) \in A(F(i)) \iff (u_{i+1}, v_{i+1}) \in A(F(i+1) \sqcup F'(i+1))$,

$$(u_i^{1 i_2 \cdots i_{n-1}}, u_i^{1 i_2 \cdots i_n}) \in A(F(i)) \iff (v_{i+1}^{i_2 \cdots i_{n-1}}, v_{i+1}^{i_2 \cdots i_n}) \in A(F(i+1) \sqcup F'(i+1)),$$

and for $i_1 \geq 2$,

$$(u_i^{i_1 i_2 \cdots i_{n-1}}, u_i^{i_1 i_2 \cdots i_n}) \in A(F(i)) \iff (u_{i+1}^{(i_1-1)i_2 \cdots i_{n-1}}, u_{i+1}^{(i_1-1)i_2 \cdots i_n}) \in A(F(i+1) \sqcup F'(i+1)).$$

Hence g is an isomorphism. $\square$

According to Lemma 6, it is straightforward to verify $|V(F(i))| = 2^{t-1-i}$, $1 \leq i \leq t-1$.

Definition 8 Let D be a digraph, $\{u, v, w\} \subseteq V(D)$, (u, v, w) is a *directed triple* of D if there is a directed path $P(u, v, w)$ from u to w , and contains v as an inner vertex. Let $S \subseteq V(D)$, S is a *3-cut set* of D if there is no directed triple of D in $V(D) - S$.

Lemma 7 $t \geq 3$, S is 3-cut set of F_t , then $|S| \geq 2^{t-3} - 1$, and there exists a 3-cut set such that ' $=$ ' holds.

proof If $t = 3$, then the length of the longest directed path of F_3 is 1. So the directed triple is $\emptyset$. Hence the proposition comes true.

Let $t \geq 4$, and suppose that S is a 3-cut set of F_t , then $S \cap V(F(i)) = S_i$ is a 3-cut set of $F(i)$, and $S = \bigcup_{i=1}^{t-1} S_i$. Assume that S_i is the smallest 3-cut set of $F(i)$, it is easy to see that we need to confirm $|S_i| = 2^{t-i-3}$, $1 \leq i \leq t-3$.

Let $S_i = \{u_i^{i_1 i_2 \dots i_n} : i_1 + i_2 + \dots + i_n + i \leq t - 3\}$, we will prove S_i is the smallest 3-cut set of $F(i)$ by induction on i .

For $i = t - 3$, $S_{t-3} = \{u_{t-3}\}$ is the smallest 3-cut set of $F(t - 3)$, the proposition holds. For i , $1 \leq i \leq t - 4$, hypothesize the conclusion holds for $i + 1$. It means that $S_{i+1} = \{u_{i+1}^{i_1 i_2 \dots i_n} : i_1 + i_2 + \dots + i_n + (i + 1) \leq t - 3\}$ is the smallest 3-cut set of $F(i + 1)$.

In the case of $F(i)$, by Lemma 6, $F(i) \cong F(i + 1) \sqcup F'(i + 1)$. Since $F(i + 1) \cong F'(i + 1)$, as the image of S_{i+1} , S'_{i+1} is the smallest 3-cut set of $F'(i + 1)$. So $S_{i+1} \cup S'_{i+1}$ is a 3-cut set of $F(i + 1) \sqcup F'(i + 1)$. By the mapping g in Lemma 6, the preimage of $S_{i+1} \cup S'_{i+1}$ is $\{u_i^{1 i_1 i_2 \dots i_n}, u_i^{(i_1+1) i_2 \dots i_n} : 1 + i_1 + i_2 + \dots + i_n + i \leq t - 3\} = S_i$, which is a 3-cut set of $F(i)$. $|S_i| = 2^{t-i-3}$ follows by induction.

If S_i is not the smallest 3-cut set of $F(i)$, then there is a 3-cut set of $F(i)$, denoted as T , such that $|T| = |g(T)| < |S_i| = 2^{t-i-3}$. Since $g(T)$ is a 3-cut set of $F(i + 1) \sqcup F'(i + 1)$, and $g(T) = (g(T) \cap V(F(i + 1))) \cup (g(T) \cap V(F'(i + 1)))$, assume $|g(T) \cap V(F(i + 1))| \leq 2^{t-(i+1)-3} - 1 < |S_{i+1}|$, contradicting to S_{i+1} is the smallest 3-cut set of $F(i + 1)$.

Hence $S = \bigcup_{i=0}^{k-1} S_i$ is the smallest 3-cut set of F_t satisfying $|S| = 2^{t-3} - 1$. $\square$

Corollary 1 Let $t \geq 4$, $U \subseteq V(F_t^\circ)$, if $|U| > 3 \cdot 2^{t-3}$, then there exists a directed triple in U .

proof Assume to the contrary that there is no directed triple in U , then $R_t(M^t(G)) - \{u_t\} - U$ is 3-cut set, and $|R_t(M^t(G)) - \{u_t\} - U| < 2^{t-1} - 1 - 3 \cdot 2^{t-3} = 2^{t-3} - 1$, contradicts to $F_t^\circ \cong F_t$ and lemma 7. $\square$

3 Main result

It is straightforward to verify the following lemma by definition of $M^t(G)$.

Lemma 8 Let v_1, v_2 be two t th roots of $M^t(K_n)$, and there is a directed path $P(v_1, v_2)$ from v_1 to v_2 in F_t° , then $N(\{v_2, h^{-1}(v_2)\}) \cap T(V(K_n)) \subseteq N(\{v_1, h^{-1}(v_1)\})$, $N(v_2) \cap T(V(K_n)) \subseteq N(v_1)$.

Lemma 9 Let $t \geq 3$, $\chi_c(M^t(K_n)) = \frac{k}{2}$, and a $(k, 2)$ -coloring satisfy Lemma 3. If there exists a set of 3 t th roots $\{v_1, v_2, v_3\}$ such that $C(v_i) \subseteq \delta(v_i) \cap R(M^t(K_n)) \subseteq \{v_i, h^{-1}(v_i)\}$, $1 \leq i \leq 3$, then (v_1, v_2, v_3) is not the directed triple of F_t° .

proof Suppose on the contrary that (v_1, v_2, v_3) is a directed triple of F_t° , and $C(v_i) \subseteq \{v_i, h^{-1}(v_i)\}$, $v_i \in X_{s_i}$, $1 \leq i \leq 3$. It is sufficient to consider two cases as follows:

Case 1. $h^{-1}(v_3) \notin C(v_3)$. According to Lemma 3, Lemma 8 and the assumption of this lemma, we get $h^{-1}(v_3) \in \delta(u_t)$, $N(v_3) \cap \delta(v_2) = \emptyset$. Hence, we can dye v_3 color s_2 , and obtain a new $(k, 2)$ -coloring $(Y_0, Y_1, \dots, Y_{k-1})$ such that $Y_{s_2} = X_{s_2} \cup \{v_3\}$, $Y_{s_3} = \emptyset$ and $Y_i = X_i, i \neq s_2, s_3$, a

contradiction to Lemma 1.

Case 2. $h^{-1}(v_3) \in C(v_3)$. Suppose that $h^{-1}(v_2) \in C(v_2)$, by Lemma 8 and the assumption of this lemma, we get $N(v_3, h^{-1}(v_3)) \cap \delta(v_2, h^{-1}(v_2)) = \emptyset$. Hence, we can dye $v_3, h^{-1}(v_3)$ color s_2 and obtain a new $(k, 2)$ -coloring $(Y_0, Y_1, \dots, Y_{k-1})$ such that $Y_{s_3} = \emptyset$, a contradiction. So $h^{-1}(v_2) \notin C(v_2)$, then we can dye v_2 color of v_1 and also obtain a new $(k, 2)$ -coloring $(Y_0, Y_1, \dots, Y_{k-1})$ such that $Y_{s_2} = \emptyset$, a contradiction. $\square$

Lemma 10 Let G be a simple graph with $V(G) = \{x_1, x_2, \dots, x_n\}$. The vertex set of $M(G)$ is $V(M(G)) = \{x_1, x_2, \dots, x_n\} \cup \{x'_1, x'_2, \dots, x'_n\} \cup \{u\}$. If $M(G)$ has (k, d) -partition, then there is a (k, d) -partition of $M(G)$ such that

1. $X_0 = \{u\}$;
2. For some i , $d \leq i \leq k - d, \forall x \in V(G)$, we have $\{x, x'\} \subseteq X_i$ if $x \in X_i$.

proof For any vertex $v \in V(M(G)) - \{u\}$, v' is the twin of v .

By Lemma 3, there exists a (k, d) -partition satisfying 2 and $u \in X_0$. If $x \neq u, x \in X_0$, and $x' \in X_i, d \leq i \leq k - d$, then $x \in V(G)$. We can dye x color i . Otherwise, there exists $y \in \delta(x')$ such that $x'y \notin E(M(G)), xy \in E(M(G))$. So $y \notin V(G) \cup \{u\}, x'y' \in E(M(G))$. Hence y' can't be dyed any color, this is impossible. Hence, $x \notin X_0, X_0 = \{u\}$ and conclusion 1 holds as well. $\square$

Theorem 1 Let $t \geq 4, n \geq \frac{11}{12}2^{t-1} + 2t + \frac{1}{3}$, then $\chi_c(M^t(K_n)) = \chi(M^t(K_n)) = n + t$.

proof Suppose that $\chi_c(M^t(K_n)) = \frac{k}{d}, d \neq 1$, by Lemma 2

$$\begin{aligned} d-1 &\leq \frac{2^t-2}{n-3} \leq \frac{2^t-2}{\frac{11}{12}2^{t-1}+2t-\frac{8}{3}} \leq 3 + \frac{2^t-2-3(\frac{11}{12}2^{t-1}+2t-\frac{8}{3})}{\frac{11}{12}2^{t-1}+2t-\frac{8}{3}} \\ &= 3 - \frac{3 \cdot 2^{t-3} + 6t - 6}{\frac{11}{12}2^{t-1}+2t-\frac{8}{3}} < 3 \end{aligned}$$

holds for $t \geq 3$. So $d \leq 3$.

If $d = 3$, there exists a $(3n + 3t - i, 3)$ -coloring of $M^t(K_n)$, $i = 1$ or 2 . According to box principle, there are at least $5n - (3n + 3t - i) = 2n - 3t + i$ color classes in the intersection sets of d -fields of different initial vertices. By Lemma 10, there are at most 3 such color classes in $\delta(u_t)$. So $V(M^t(K_n)) - \delta(u_t)$ have at least $2n - 3t + i - 3$ such color classes in which contains at least one vertex in $R_t(M^t(G)) - \{u_t\}$ respectively. Thus $2n - 3t + i - 3 \leq 2^{t-1} - 1$, i.e.,

$$n \leq 2^{t-2} + \frac{3}{2}t + \frac{2-i}{2} \leq 2^{t-2} + \frac{3}{2}t + \frac{1}{2}.$$

If $t \geq 4$, then

$$\frac{11}{12}2^{t-1} + 2t + \frac{1}{3} - (2^{t-2} + \frac{3}{2}t + \frac{1}{2}) = \frac{5}{6}2^{t-2} + \frac{1}{2}t - \frac{1}{6} > 0.$$

So

$$n \geq \frac{11}{12}2^{t-1} + 2t + \frac{1}{3} > 2^{t-2} + \frac{3}{2}t + \frac{1}{2} \geq n,$$

a contradiction.

Suppose that $d = 2$, then $M^t(K_n)$ has a $(2n + 2t - 1, 2)$ -coloring $(X_0, X_1, \dots, X_{2n+2t-2})$ satisfying Lemma 10. According to box principle, there are at least $3n - (2n + 2t - 1) = n - 2t + 1$ color classes in the intersection set of d -fields of different initial vertices. By Lemma 10, there are at most 1 such color class in $\delta(u_t)$. By definition 5, any of these color classes we get is in $R(M^t(K_n))$, and has at least one t th root.

Now we discuss such color classes in $V(M^t(K_n)) - \delta(u_t)$. It is easy to see that the number of these color classes is at least $n - 2t$. Suppose that the number of these color classes that contain exactly 1 t th root is y , the number of those which contain at least 2 t th roots is x and the number of t th roots which is different from $\{u_t\}$ and not in such intersection sets is z , then we have

$$x + y \geq n - 2t, \quad 2x + y + z \leq 2^{t-1} - 1.$$

Hence

$$y \geq 2n - 2^{t-1} - 4t + 1 + z, \quad z \leq 2^{t-1} + 2t - n - 1.$$

Since $d = 2$, the color classes in the intersection set of d -field of different initial vertices are mutually disjoint. Hence, in the set of y classes which contain exactly 1 t th root, the number of classes whose d -fields contain other t th roots is at most $2z$, furthermore there are at most 2 classes whose d -fields contain r th roots, $1 \leq r \leq t - 1$, X_2 and $X_{2n+2t-3}$. Hence, there are at least $y - 2z - 2$ color classes whose d -fields contain exactly 1 t th root.

By Corollary 1, if $y - 2z - 2 > 3 \cdot 2^{t-3}$, then there exists a directed triple of F_t° in the set of t th roots that contained in these $y - 2z - 2$ color classes, contradict to Lemma 9. Hence, $y - 2z - 2 \leq 3 \cdot 2^{t-3}$, we have

$$\begin{aligned} & (2n - 2^{t-1} - 4t + 1 + z) - 2z - 2 \leq y - 2z - 2 \leq 3 \cdot 2^{t-3} \\ \implies & 2n - 2^{t-1} - 4t - 1 - z \leq 3 \cdot 2^{t-3} \\ \implies & 2n - 2^{t-1} - 4t - 1 - (2^{t-1} + 2t - 1 - n) \leq 3 \cdot 2^{t-3} \\ \implies & 3n \leq 11 \cdot 2^{t-3} + 6t \\ \implies & n \leq \frac{11}{12} \cdot 2^{t-1} + 2t, \end{aligned}$$

contradict to $n \geq \frac{11}{12}2^{t-1} + 2t + \frac{1}{3}$. So $d \neq 2$.

In other words, if $n \geq \frac{11}{12}2^{t-1} + 2t + \frac{1}{3}$, then $d = 1$, $\chi_c(M^t(K_n)) = \chi(M^t(K_n)) = n + t$. $\square$

References

- [1] J. A. Bondy and U. S. R. Murty, Graph Theory with Applications, Macmillan Press, London, 1976.
- [2] J. A. Bondy and P. Hell, A Note on the Star Chromatic Number, J. Graph Theory, 1990,14:479-482.
- [3] G. J. Chang, L. Huang and X. Zhu, Circular Chromatic Number of Mycielski's Graphs, Discrete mathematics, 1999,205:23-37.
- [4] Genghua Fan, Circular Chromatic Number and Mycielski Graphs, Combinatorica,2004, 24:127-135.
- [5] D. R. Guichard, Acyclic Graph Coloring and the Complexity of the Star Chromatic Number, J. Graph Theory,1993,17:129-134.
- [6] H. Hajiabolhassan and X. Zhu, Circular Chromatic Number and Mycielski Construction, J. Graph Theory, 2003,44:106-115.
- [7] F. Harary, Graph Theory, Addison-Wesley, New York, 1969.
- [8] L. Huang and G. J. Chang, The circular Chromatic Number of the Mycielskian of G_k^d , J. Graph Theory, 1999,32:63-74.
- [9] M. Kubale, Graph Colorings, American Mathematical Society, Providence,Rhode island,2004.
- [10] D. D. F. Liu, Circular Chromatic Number for Iterated Mycielski Graphs, Discrete mathematics, 2004,285:335-340.
- [11] Hongmei Liu, On a Conjecture of Circular Chromatic Number of Mycielski Graphs, Journal of Mathematical, 2006,26:602-608.
- [12] G.Simonyi and G.Tardos,Local chromatic number,Ky Fan's theorem and circular colorings,Combinatorica 2006,26:587-626.
- [13] A. Vince, Star Chromatic Number, J. Graph Theory, 1988,12:551-559.
- [14] X. Zhu, Star Chromatic Number and Products of Graphs, J. Graph Theory,1992,16:557-569.
- [15] X. Zhu, Circular Chromatic Number: A Survey, Discrete mathematics,2001,229:371-410.